


Sample Name:

Poly(methyl methacrylate-*co*_(random)-n-butyl methacrylate)-*block*-poly(2-hydroxyethyl methacrylate)

Sample #: P10580-MMA_nBuMA_ran-b-HEMA

Structure:



Composition:

$M_n \times 10^3$ (g/mol)	22.0-16.5
M_w/M_n	1.19
Molar ratio MMA : nBuMA	54 : 46 (mol/mol)
Weight ratio MMA : nBuMA	26 : 31 : 43 (wt%)
T_g (MMA _n BuMA)	65 °C
T_g (HEMA)	112 °C

Synthesis Procedure:

Poly([methyl methacrylate-*co*-n-butyl methacrylate]-*b*-2-hydroxyethyl methacrylate) block copolymer was synthesized by living anionic polymerization. First, methyl methacrylate (MMA) and n-butyl methacrylate (n-BuMA) were co-polymerized; and then 2-[trimethylsilyloxy]ethyl methacrylate (hydroxyprotected HEMA monomer) was added. The obtained block copolymer was precipitated in acidic methanol solution to deprotect the hydroxyl group.

Solubility: The polymer is soluble in THF, DMF.


Characterization:

The polymer composition was determined by ¹H NMR. MMA:nBuMA molar ratio was calculated by comparing the integration of the -OCH₂- protons of nBuMA (at δ = 3.9 ppm) to the integration of methoxy group of MMA (at δ = 3.6 ppm). Molecular weight of the second (HEMA) block was calculated by comparing the integration of -OCH₂- protons of HEMATMS to the integration of methoxy group of MMA and using SEC data for the first (MMA_nBuMA) block.


The average molecular weight and polydispersity index were determined by size exclusion chromatography (SEC). For SEC analysis, the MMA_nBuMA-b-HEMA block copolymer can be treated with acetic anhydride in presence of pyridine to convert the hydroxy-groups to acetate groups.

Thermal analysis of the sample was done on a TA Q100 differential scanning calorimeter (DSC) at a heating rate of 10°C/min. The glass transition temperature (T_g) was determined as a midpoint of step change in heat flow curve for the second heating scan.

¹H NMR of MMA_nBuMA_ran [first block]:




¹H NMR of MMA_nBuMA_ran-b-HEMA diblock copolymer:




SEC of MMA_nBuMA_ran [first block]:

Sample ID: P10580-I-MMA_nBuMA

Concentration (mg/mL)	6.366
Sample dn/dc (mL/g)	0.080
Method File	PS8K-Oct-0000.vcm
Column Set	3x PL 1113-6300
System	System 1




Sample	Mn (Da)	Mw (Da)	Mp (Da)	Mw/Mn	IV (dL/g)
P10580-1_2011-11-21_13:59:25_01.vdi	21,864	24,001	22,443	1.098	0.1836



SEC of MMA_nBuMA_ran and MMA_nBuMA_ran-b-HEMATMS:

P10580-MMA_nBuMA_ranHEMA



Size exclusion chromatography of
1. Poly MMA_nBuMA_ran: Mn 2,000 Mw: 24,000 Mw/Mn 1.09
Poly MMA_nBuMA_ran-b-HEMATMS: Mn 22,000 -b-25,600 Mw/Mn 1.19
After Deprotection : MMA_nBuMA_ran-b-HEMA Mn 22,000-b-16,500
It shows no micellization in THF